

Fig. 2. Isothermal compressibility  $\chi_T$  for Kr

○ Stewart (13); □ Urvas et al. (10); ● this work

on similar crystals revealed that the specimens consisted of grains with diameters of approximately 5 mm (5). After the crystals were grown, they were transferred from the growing tube to the dilatometer chamber.

For the compressibility measurements a pressure from 1 to 20 bar was applied to the specimen by gaseous or liquid helium. The change of sample length was independent of whether the pressure was being raised or lowered; i. e., no hysteresis




Fig. 3. Comparison of the experimental volume thermal expansion coefficient for Kr with theory (1). MLJ (12, 6) means (12, 6)-Mie-Lennard-Jones potential; qh: quasiharmonic, anh: anharmonic approximation; a: measured with X-rays (9); b: bulk expansivity measured in this work

— x — qh. MLJ (12, 6),      - - - - - anh. MLJ (12, 6),  
 - · - · - - anh. MLJ (13, 6),      ——— experiment

Fig. 4.

isotherm

theory (

(9, 10);

ed in th

tice and

— x —

- · - · -

— · - -

—

effect c

pansivi

M

in the t

(6 to 8)

(Fig. 1

In

and the

et al. (

from th

of our

al. (9)

therma

by Lose

F

compre

tions (1

T