

COHE-LW 75-0407

DETERMINATION OF HIGH-TEMPERATURE TRANSITION IN CALCITE TO 5 KBAR BY DIFFERENTIAL THERMAL ANALYSIS IN HYDROSTATIC APPARATUS

LEWIS H. COHEN AND WILLIAM KLEMENT, JR.

Department of Geological Sciences, University of California, Riverside, California 92502;
Department of Engineering and Applied Science, University of California, Los Angeles, California 90024

ABSTRACT

The high-temperature transition in calcite, located by extrapolation near 985° at 1 bar, has been determined to 5 kbar by differential thermal analysis in hydrostatic apparatus. The transition temperature increases with pressure at the rate of $3.0 \pm 0.3^\circ\text{C}/\text{kbar}$, in contrast to the recent hypotheses which project a decrease.

INTRODUCTION

In a thermal analysis study of calcite, Boeke (1912) reported a subtle anomaly near $970 \pm 5^\circ\text{C}$. Existence of the anomaly was corroborated by Eitel (1923), who placed it near $975\text{--}980^\circ\text{C}$ under ~ 0.1 kbar CO_2 pressure. However, Smyth and Adams (1923) could not verify the anomaly in their careful differential thermal analyses. Following Jamieson's (1957) suggestion, Boettcher and Wyllie (1967) strongly inferred that this anomaly in calcite corresponded to a transition whose temperature decreased with pressure to near 480°C and 9.4 kbar in order to explain a kink in the trajectory of the calcite-aragonite boundary there. Goldsmith and Newton (1969) confirmed the kink discovered by Boettcher and Wyllie (1967).

¹ Manuscript received February 21, 1973. [JOURNAL OF GEOLOGY, 1973, Vol. 81, p. 724-727] © 1973 by the University of Chicago. All rights reserved.

and also unsuccessfully sought to confirm the anomaly in calcite by several methods, including differential thermal analysis (DTA) in piston-cylinder apparatus. Zimmernann (1971) apparently located no change in slope within his experimental range, but Johannes and Puhar (1971), in subsequent work, found a gradual curve in the trajectory of the transition between 300° and 500°C . The present DTA investigation in hydrostatic apparatus has established the trajectory of the thermal anomaly in calcite, which does not correspond to the trajectory postulated by Jamieson (1957) and others.

EXPERIMENTS AND RESULTS

AR primary standard CaCO_3 , obtained from Mallinckrodt and verified by X-ray diffraction to be calcite, was dried at $\sim 200^\circ\text{C}$ for 1 day under 0.5 atm of air.